NC Violent Death Reporting System

SUICIDE AMONG NON-HISPANIC AFRICAN AMERICAN YOUTH, AGES 10-24, IN **NORTH CAROLINA: 2014-2023**


The NC-VDRS is a CDC-funded statewide surveillance system that collects detailed information on violent deaths that occur in North Carolina including: homicide, suicide, unintentional firearm deaths, legal intervention, and deaths of undetermined intent. NC-VDRS triangulates information from death certificates, medical examiner reports, and law enforcement reports to understand and aid in the prevention of violent deaths. For more information, please visit https://injuryfreenc.dph.ncdhhs.gov/About/ncVDRS.htm.

Non-Hispanic African American Youth Suicide in Number of North Carolina, by Sex and Age: **Suicides** 45 ■ Males ■ Females 40 35 30 25 20 15 10 5 10 12 13 14 15 16 17 18 20 23 Age

- From 2014 to 2023, 1,963 non-Hispanic (NH) African American North Carolina residents ages 10 to 24 died as a result of violence. Of these violent deaths, 408 (20.8%) were suicides.
- Males consistently had a higher number of suicides than females, regardless of age.
- The number of suicide deaths peaked for NH African American males at age 23 (40 suicides) and for NH African American females at age 16 (10 suicides).

- Overall, the most common method of suicide among NH African American youth was firearms (56.6%), followed by hanging, strangulation, suffocation (30.9%) and poisoning (5.4%).
- Only 7.2% of suicides involved other methods.

Method of Suicide Among NH African American Youth: NC-VDRS, 2014-2023




*Other includes falls, drowning, motor vehicle, fire/burns and other causes of suicide.

If you or someone you know needs support now,



call or text 988 or chat 988lifeline.org

Precipitating Circumstances of NH African American Youth Suicide by Sex: NC-VDRS, 2014-2023



- Fifty-four percent (53.6%) of female and 38.7% of male suicide victims were characterized as having a current mental health problem when they died by suicide.
- A similar trend was seen for mental health treatment. Females (33.3%) were more likely than males (19.2%) to be receiving treatment for a current mental health problem at the time of suicide.
- Twenty-nine percent (28.6%) of female and 28.2% of male suicide victims had an intimate partner problem.
- Forty-eight percent (47.6%) of female and 47.4% of male suicide victims had experienced a recent or imminent crisis.
- Twenty-three percent (22.6%) of female and 26.8% of male suicide victims had disclosed their suicide intent to someone else before they died.
- Females (31.0%) were more likely to leave a suicide note than males (19.5%).

More infromation on suicide prevention efforts can be found at:

State Resource Partners

Division of Mental Health,

Developmental Disabilities and Substance Use Services

https://www.ncdhhs.gov/divisions/mhddsus

North Carolina Office of the Chief Medical Examiner

https://www.ocme.dhhs.nc.gov/

Injury and Violence Prevention Branch Suicide Prevention Website

https://injuryfreenc.dph.ncdhhs.gov/preventionResources/Suicide.htm

Division of Child and Family Wellbeing

https://www.ncdhhs.gov/divisions/division-child-and-family-well-being

National Resources The Suicide Prevention Resource Center http://www.sprc.org/

The American Foundation for Suicide Prevention

https://afsp.org/

The National Suicide Prevention Lifeline

call or text 988

or chat 988lifeline.org

The North Carolina Violent Death Reporting System is supported by Cooperative Agreement 5 NU17CE924955-02-00 from the Centers for Disease Control and Prevention (CDC).





NCDHHS Division of Public Health / www.dph.ncdhhs.gov / Injury Epidemiology & Surveillance Unit/ 919-707-5425 NC Violent Death Reporting System / 919-707-5432 State of North Carolina / Department of Health and Human Services / www.ncdhhs.gov

NCDHHS is an equal opportunity employer and provider. 2023 PROVISIONAL DATA 6/20/2025

Please see the NC-VDRS 2023 Annual Report for additional data and technical information:

 $\underline{https://injuryfreenc.dph.ncdhhs.gov/DataS}urveillance/ViolentDeathData.htm$

^{*}Circumstances are known for 94% of males (n=302) and 96% of females (n=84)