
NC Violent Death Reporting System

SUICIDE AMONG OLDER ADULTS IN NORTH CAROLINA, 2019 - 2023

The North Carolina Violent Death Reporting System (NC-VDRS) is a CDC-funded statewide surveillance system that collects detailed information on violent deaths that occur in North Carolina including: homicide, suicide, unintentional firearm deaths, legal intervention, and deaths of undetermined intent. NC-VDRS uses information from death certificates, medical examiner reports, and law enforcement reports to understand and aid in the prevention of violent deaths. For more information, please visit https://injuryfreenc.dph.ncdhhs.gov/About/ncVDRS.htm.

Older Adult Suicide Rates* by Age Group: NC-VDRS, 2019 - 2023




- From 2019 to 2023, 1,872 North Carolina residents ages 65 and older died as a result of violence. Of these violent deaths, 1,568 were suicide (83.8%).
- The suicide rate among older adults was 17.4 suicides per 100,000 population.
- Eighty-two percent (82.3%) of suicides among older adults were among males.
- The suicide rate peaked among males 85 and older, and the suicide rate was consistently higher for males compared to females for all age groups.
- Among females, the suicide rate peaked among those ages 65-74.
- Ninety-three (92.4%) percent of older adult suicide victims were identified as non-Hispanic (NH) White, 5.1 percent were NH Black, and the remaining 2.5 percent were identified as NH Asian, NH American Indian, or Hispanic.
- Overall, the most common method of suicide among older adults was firearms (76.6%), followed by poisoning (10.9%) and hanging/strangulation/suffocation (8.4%).
- Four percent (4.1%) of suicides involved a method other than firearms, poisoning, or hanging/strangulation/suffocation.

If you or someone you know needs support now,




call or text 988 or chat 988lifeline.org

Elder Suicide Method of Death: NC-VDRS, 2019 - 2023



*Other methods include: sharp instrument, fire/burns, motor vehicle and other.

Older Adult Suicide Circumstances*: NC-VDRS, 2019 - 2023



- Over one-third (33.9%) of older adult male suicide victims with circumstance information were characterized as having a current depressed mood at the time of death compared to 35.6% of females.
- Sixty-seven percent (66.7%) of older adult female and 51.3% of older adult male suicide victims had a current mental health problem when they died by suicide.
- Twenty-one percent (20.6%) of older adult females had a history of prior suicide attempts, whereas only 7.1% of older adult males had a history of prior suicide attempts.
- Sixty-two percent (62.0%) of males and 49.1% of females had a physical health problem.

More information on suicide prevention efforts can be found at:

State Resource Partners

Division of Mental Health,

Developmental Disabilities and Substance Use Services

https://www.ncdhhs.gov/divisions/mhddsus

Office of the Chief Medical Examiner

https://www.ocme.dhhs.nc.gov/

North Carolina Center for Aging and Health

https://www.med.unc.edu/aging/

Division of Aging

https://www.ncdhhs.gov/divisions/daas

National Resources
The Suicide Prevention Resource Center

http://www.sprc.org/

The American Foundation for Suicide Prevention

https://afsp.org/

The National Suicide Prevention Lifeline

call or text 988 or chat 988lifeline.org

The North Carolina Violent Death Reporting System is supported by Cooperative Agreement 5 NU17CE924955-02-00 from the Centers for Disease Control and Prevention (CDC).





NCDHHS Division of Public Health / www.publichealth.nc.gov / Injury Epidemiology & Surveillance Unit/ 919-707-5425

NC Violent Death Reporting System / 919-707-5432

NA VIOLENT DEATH

State of North Carolina / Department of Health and Human Services / www.ncdbhs.gov

State of North Carolina / Department of Health and Human Services / www.ncdhhs.gov NCDHHS is an equal opportunity employer and provider. FINAL DATA 6/16/2025

Please see the NC-VDRS 2023 Annual Report for additional data and technical information:

https://injuryfreenc.dph.ncdhhs.gov/DataSurveillance/ViolentDeathData.htm

^{*}Circumstances are known for 98% of males (n=1258) and 96% of females (n=267).