





Gonorrhea in North Carolina, 2024

Gonorrhea may be declining in North Carolina.

Gonorrhea cases increased **49%** compared to 2014, but declined 16% compared to 2023.




A little more than half of people with gonorrhea were **men (58%)**. **Gonorrhea decreased among both men and women in 2024.**



Disparities by income level are particularly large for gonorrhea

Gonorrhea rates are highest among people living in the most impoverished neighborhoods.

People living in impoverished areas often have less access to resources, including health resources. Delayed treatment can increase the potential for transmission to others.



*Estimates of people living below the poverty line within a census tract and all population estimates obtained from the American Community Survey, 2019-2023, five-year estimate

Proportion of Census Tract Living in Poverty

Most people with gonorrhea did not identify as Hispanic/Latine; however, there was a greater decrease in gonorrhea among non-Hispanic/non-Latine people in 2024 (15% decrease) than among Hispanic/Latine people (6% decrease).

In 2024, 14% of men and <1% of women diagnosed with gonorrhea were coinfectd with HIV. Clinicians should offer HIV testing to all patients diagnosed with gonorrhea and discuss HIV pre-exposure prophylaxis (PrEP) with those who test HIV-negative.

More information: <https://epi.dph.ncdhhs.gov/cd/stds/annualrpts>

*2020 data should be treated with caution due to reduced availability of testing caused by the COVID-19 pandemic.

Data Source: North Carolina Electronic Disease Surveillance System (NC EDSS) (data as of July 1, 2025) and enhanced HIV/AIDS Reporting System (eHARS) (data as of July 2025)

Contact Us:
North Carolina Department of Health and Human Services
Division of Public Health
Communicable Disease Branch
Phone: (919) 733-3419

Mailing Address: Communicable Disease Branch Epidemiology Section
1902 Mail Service Center
Raleigh, NC 27699-1902
Prepared by the HIV/STD/Hepatitis Surveillance Unit
Created on 9/10/2025




Gonorrhea in North Carolina, 2024



What is North Carolina doing about gonorrhea?

- North Carolina funds gonorrhea screening for all women who are seen in publicly funded health care facilities, such as local health departments and family planning settings.
- Nurses in the North Carolina Division of Public Health Technical Assistance and Training Program work with county staff to ensure correct treatment of gonorrhea.
- North Carolina is participating in a nation-wide program to identify drug-resistant gonorrhea now called Combatting Antimicrobial Resistant Gonorrhea and Other STIs (CARGOS). For more information: <https://www.cdc.gov/sti/php/projects/cargos.html>.
- Doxy PEP is a new approach to addressing STI prevention and has demonstrated benefit in reducing STI incidence (e.g., syphilis, chlamydia, and gonorrhea). For more information: [Preventing STIs with Doxy PEP](#).

Treatment for Gonorrhea



- First-line Recommendation
- Second-line Recommendation
- Other Treatment
- No treatment reported

2024 Gonorrhea Treatment

- Correctly treating gonorrhea is very important to prevent the development of drug resistance.
- CDC [STI Treatment Guidelines](#) for gonorrhea include:
 - Recommended first-line regimen: Ceftriaxone (500 mg intramuscular (IM) in a single dose for individuals <150 kg or 1 gm for individuals ≥150 kg).
 - Second-line recommendations: Cefixime (800 mg orally in a single dose) OR gentamicin (240 mg IM in a single dose) plus azithromycin (2gm orally in a single dose).

Most people with gonorrhea are receiving the first-line recommendation.

What CLINICIANS can do

- Always ask patients about their sexual activity and test those that are sexually active at all sites of exposure (e.g., pharynx, urethra, and rectum); retest 3 months after treatment and refer sex partners for evaluation and treatment.
- Provide treatment consistent with the CDC guidelines (<https://www.cdc.gov/std/treatment-guidelines/gonorrhea.htm>).
- Suspected cases of gonorrhea [treatment failure](#) should be reported immediately to [your local health department](#) or NC Division of Public Health (919-733-3419).
- For other resources, visit the National Coalition for Sexual Health compendium: [Compendium of Sexual Reproductive Health Resources for Healthcare Providers](#).

What YOU can do

If you have a gonorrhea infection, ensure that you and your partners get treatment, and retested after 3 months. Untreated gonorrhea can lead to increased risk for HIV and pelvic inflammatory disease (PID).

Data Source: North Carolina Electronic Disease Surveillance System (NC EDSS) (data as of July 1, 2025) and enhanced HIV/AIDS Reporting System (eHARS) (data as of July 2025).

State of North Carolina • Josh Stein, Governor
NC DHHS • Devdutta Sangvai, MD, Secretary
NC DPH • Kelly Kimple, MPH, MD, Director Division of Public Health
HIV/STD/Hepatitis Surveillance Unit • Erika Samoff, MPH, PhD

www.ncdhhs.gov • www.publichealth.nc.gov
NC DHHS is an equal opportunity employer and provider
Prepared by the HIV/STD/Hepatitis Surveillance Unit
Created on 9/10/2025